



Transmission of  
Viruses in Droplets  
and Aerosols

LINSEY C. MARR

*CHARLES P. LUNSFORD PROFESSOR  
CIVIL AND ENVIRONMENTAL ENGINEERING  
VIRGINIA TECH*

26 MARCH 2020



# Topics


1. Respiratory viruses
2. Transmission modes
3. Size distributions and evaporation
4. Virus aerosol dynamics
5. Impact of temperature and humidity
6. Masks
7. SARS-CoV-2

## Viruses that infect the upper respiratory tract

Rhinovirus  
Coronavirus  
Influenza virus  
Parainfluenza virus  
Respiratory Syncytial virus  
Herpesvirus  
Adenovirus  
Bocavirus  
Coxsackivirus

## Viruses that infect the lower respiratory tract

Influenza virus  
Parainfluenza virus  
Respiratory Syncytial virus  
Adenovirus  
Bocavirus  
Metapneumovirus



# Virus Size







Image courtesy of the U.S. EPA




influenza  
0.1  $\mu\text{m}$



SARS-CoV-2  
0.12  $\mu\text{m}$





rhinovirus  
0.03  $\mu\text{m}$



adenovirus  
0.1  $\mu\text{m}$

# Size Matters


- Airborne virus is not naked!




- Size determines
  - Lifetime in the atmosphere
  - Where it deposits in the respiratory system



# Modes of Transmission




direct contact




indirect contact

Defined as  $>5 \mu\text{m}$   
and happening at  
close-range only  
( $<2 \text{ m}$ )




large droplets



Defined as  $<5 \mu\text{m}$   
and happening  
mainly at long-  
range ( $>2 \text{ m}$ )

aerosols

# Reality is More Complicated



Linsey Marr, Virginia Tech, March 2020


Droplets that are expelled into air can be inhaled, land on people's mucus membranes, or deposit onto surfaces, where someone can touch them or they can be resuspended into air.

What size are these droplets?




# Size Distributions: Breathing

Measured by  
aerodynamic particle  
sizer over the range  
0.3-20  $\mu\text{m}$




# Size Distributions: Speaking

Measured by  
aerodynamic particle  
sizer over the range  
0.3-20  $\mu\text{m}$



# Size Distributions: Coughing


Measured by  
aerodynamic particle  
sizer over the range  
0.3-20  $\mu\text{m}$





# Corrected Size Distributions

Inferred number concentration in upper respiratory tract



Breathing, talking, and coughing release droplets that range from submicron to millimeter in size.



What size droplets carry viruses?

# Virus Detection Methods

## 1. Total virus

- Number of genome copies (GC) determined by molecular techniques (quantitative polymerase chain reaction, qPCR)
- Reflects number of viruses with intact DNA or RNA
- Does NOT indicate whether virus is infectious or not

## AN INFLUENZA VIRUS




RNA is wrapped around the ribonucleoprotein


# Virus Detection Methods

## 2. Infectious virus

- Number of viruses that are able to infect cells determined by culture (growing)
- PFU = plaque forming units, number of viruses capable of forming plaques on host cells, focus forming units (FFU) are related
- $TCID_{50}$  = median tissue culture infectious dose, concentration at which half of cells are infected after being exposed to the sample




# Relationship Between the Two Methods for Flu Virus



There is a weak, but significant, correlation between virus RNA copies and infectious virus.





# Amount of Flu Virus in Coarse vs. Fine Droplets (Particles) in Exhaled Breath



**Figure 1. Influenza virus copy number in aerosol particles exhaled by patients with and without wearing of an ear-loop surgical mask.** Counts below the limit of detection are represented as 0.5 on the log scale. doi:10.1371/journal.ppat.1003205.g001

# Flu Virus in Droplets (Aerosols)



Linsey Marr, Virginia Tech, March 2020


30-min sample

recite alphabet at 5, 15, 25 min


The majority of flu virus (RNA copies) is found in fine ( $<5 \mu\text{m}$ ), rather than coarse ( $>5 \mu\text{m}$ ), droplets/aerosols.

How do these droplets move around the indoor environment?


# Settling Velocity and Time



# Humidity Controls Final Size



# Virus Dynamics in Indoor Air



$$\frac{dC_d}{dt} = -\left(\frac{v}{H} + \lambda + k\right)C_d$$

settling     ventilation (air-exchange)     inactivation

concentration of infectious virus in aerosols of diameter  $d$


- Settling velocity  $v$  depends on diameter  $d$
- Diameter depends on RH
- Inactivation rate  $k$  depends on RH



relative humidity (RH)




# Virus Viability vs. RH



# Virus-Aerosols From a Cough

$\lambda = 1$  ACH at RH = 50%




There is a size shift due to loss of larger droplets by gravitational settling.

Linsey Marr, Virginia Tech, March 2020





# Infectious Concentrations vs. RH




Concentrations are higher at lower RH mainly because lab-determined inactivation rate is lower.


# RH and Removal Mechanisms

- Settling: main removal mechanism, efficient for large but not small droplets
- Ventilation: effective for all sizes, important in public places
- Inactivation: effective for all sizes, important for small droplets

**A**  $\lambda = 1$  ACH




**B**  $\lambda = 10$  ACH









Viruses can be removed from indoor air by settling, ventilation, and inactivation; some of these processes depend on humidity.

How do temperature and humidity affect transmission?

# Seasonality of the Flu








# Virus Viability

- Temperature (T): In general, viruses survive better at lower T.
- Relative humidity (RH): Many, but not all, viruses in aerosols and droplets survive best at low RH (<40%). Some survive well at very high RH (>95%).
- Indoor T and RH are key because most transmission probably occurs indoors.






# Virus Viability vs. RH



Conflicting results in the literature




# Flu Remains Viable at All RH



Virus in L-15 medium + human bronchial epithelial cell wash maintains high viability across all RHs tested.

# Respiratory Secretions Protect

## H1N1 flu virus in droplets



Viability is lower without human bronchial epithelial cell wash. Also for bacteriophage  $\Phi 6$ .

# SARS-CoV-1 in Droplets




FIGURE 2: Infectivity of SARS Coronavirus ( $10^5/10 \mu\text{L}$ ) to different temperatures at (a) >95% relative humidity, (b) >80–89%.

Dried SARS-CoV-1 on plastic decayed faster at higher temperature and faster at >95% RH than at 80-89% RH. In another study, it decayed much more quickly at 56 and 60 °C than at 4 °C.


# How Might RH Affect Transmission?

Low RH

Medium RH

Very high RH

Physics






settles in  
8 min



settles in  
1 min

Chemistry



Viruses in air and on surfaces survive better at lower temperatures. Survival varies with humidity and liquid composition.

How do masks work?

# Types of Masks



surgical mask

intended to keep the wearer from spraying droplets onto others



respirator

intended to reduce the wearer's exposure to inhaled particles


# Three Key Factors Required for a Respirator to be Effective



- ① The respirator must be put on correctly and worn during the exposure.
- ② The respirator must fit snugly against the user's face to ensure that there are no gaps between the user's skin and respirator seal.



- ③ The respirator filter must capture more than 95% of the particles from the air that passes through it.




\*If your respirator has a metal bar or a molded nose cushion, it should rest over the nose and not the chin area.



# Filtration Mechanisms


- Impaction
- Interception
- Diffusion
- Not sieving!



“collector” = single fiber in a filter



# Filter Efficiency




# N95

- Blocks at least 95% of particles of diameter  $0.3\ \mu\text{m}$
- Removal efficiency is even better for particles  $>0.3\ \mu\text{m}$  and particles  $<0.3\ \mu\text{m}$
- Capture efficiency depends on the size and density of the particle and should be the same whether the particle contains a virus or not



Linsey Marr, Virginia Tech, March 2020





# Surgical Masks and Flu Virus




Different types of surgical masks reduced the amount of infectious flu virus measured behind the mask on a manikin by an average of a factor of 6 (range 1.1-55).

What do we know about SARS-CoV-2 in droplets/aerosols?

# Epidemiological Comparison of Respiratory Viral Infections

| Disease   | Flu  | COVID-19   | SARS  | MERS  |
|---|--|--|---|---|
| Disease Causing Pathogen                          | <br>Influenza virus | <br>SARS-CoV-2 | <br>SARS-CoV | <br>MERS-CoV |
| <b>R<sub>0</sub></b><br>Basic Reproductive Number | <b>1.3</b>   | <b>2.0 - 2.5 *</b>   | <b>3</b>  | <b>0.3 - 0.8</b>  |
| <b>CFR</b><br>Case Fatality Rate                  | 0.05 - 0.1%  | ~3.4% *  | 9.6 - 11%   | 34.4%   |
| <b>Incubation Time</b>                            | 1 - 4 days   | 4 - 14 days *  | 2 - 7 days  | 6 days  |
| <b>Hospitalization Rate</b>                       | 2%   | ~19% *   | Most cases  | Most cases  |
| <b>Community Attack Rate</b>                      | 10 - 20%   | 30 - 40% *   | 10 - 60%  | 4 - 13%   |
| <b>Annual Infected (global)</b>                   | ~ 1 billion  | N/A (ongoing)  | 8098 (in 2003)  | 420   |
| <b>Annual Infected (US)</b>                       | 10 - 45 million  | N/A (ongoing)  | 8 (in 2003)   | 2 (in 2014)   |
| <b>Annual Deaths (US)</b>                         | 10,000 - 61,000  | N/A (ongoing)  | None (since 2003)   | None (since 2014)   |

\* COVID-19 data as of March 2020.



**+ positive**

table, bed rail, locker, chair, light switches, stethoscope, sink, floor, window, door and handle, toilet, **air outlet fan**

**- negative**

control panel on bed, call bell, rail in bathroom, outer rim of sink in bathroom, all anteroom, **air**

Samples in patient A's and B's rooms were negative after cleaning. These results are for patient C before cleaning.

# SARS-CoV-2 Size Distributions









Figure 1 Concentration of airborne SARS-CoV-2 RNA in different aerosol size bins

# SARS-CoV-2 Survival in Aerosols

## C Half-Life of Viable Virus


### Aerosols





# Major Unknowns

- Which transmission route is dominant: direct contact, indirect contact with contaminated objects (fomites), inhalation of aerosols, deposition of droplets?
- How are viruses inactivated in air and on surfaces?
- How much virus is released in what size aerosols at different stages of infection?
- How well does SARS-CoV-2 survive in aerosols under real-world conditions?



# Acknowledgments

Karen Kormuth  
Seema Lakdawala  
Weinan Leng  
Kaisen Lin  
AJ Prussin II  
Elankumaran Subbiah  
Eric Vejerano  
Peter Vikesland  
Haoran Wei  
Wan Yang



Center for the Environmental  
Implications of NanoTechnology

Linsey Marr, Virginia Tech, March 2020

